

JOURNAL OF DIFFERENTIAL EQUATIONS 5, 215–225 (1969)

Two Point Boundary Problems in Banach Space

KENNETH A. HEIMES

Department of Mathematics, Iowa State University, Ames, Iowa 50010

Received October 24, 1967

1. INTRODUCTION

Let X be a Banach space, $[X]$ the Banach algebra of bounded linear operators on X into X . The continuity, derivative and integral of a function $y(t)$, or $Y(t)$, from the real interval $t \geq 0$ into X , or $[X]$, is defined in terms of the norm topology. These concepts together with the necessary calculus may be found in [1], pp 58–123. It is assumed throughout that A and B are continuous mappings from $t \geq 0$ into $[X]$.

We consider below the questions of existence and uniqueness of solutions to

$$y''(t) = A(t)y'(t) + B(t)y(t) \quad (1)$$

satisfying the boundary conditions

$$y(0) = \alpha, \quad y(\tau) = \beta \quad \alpha, \beta \in X \quad (2)$$

where y is of class C^2 on the compact interval $[0, \tau]$ into X . Our main objective is to impose conditions on the operator functions A, B and the interval length $\tau > 0$ so that (1), (2) has a unique solution for every choice of boundary values α, β .

Applying an operational calculus for commutative normed rings, we give a necessary and sufficient condition for unique solvability when A and B are constant and commute. In general, however, we can only state quantitative sufficient conditions (i.e. inequalities involving norms), some of which generalize classical results for the scalar case. When X is Hilbert space, several qualitative conditions on A and B are given which make (1), (2) uniquely solvable.

If X has finite dimension, solutions to (1), (2) are unique if and only if (1), (2) is solvable for all $\alpha, \beta \in X$. ([2], p. 387) Consequently, a study of the boundary value problem reduces to determining conditions on the $n \times n$ matrices A, B and the interval length $\tau > 0$ under which the only solution to (1) vanishing at $t = 0, \tau$ is the trivial solution; i.e. disconjugacy or non-

oscillation conditions. Hartman and Wintner have given disconjugacy criteria in [3] when (1) is non-self-adjoint. For the self-adjoint case, sufficient conditions for disconjugacy have been given by Barrett [4] and Reid [5]. The related questions of oscillation or non-oscillation for large t for the system (1) and/or the corresponding matrix system are considered in [4]–[10].

2. PRELIMINARIES

Let Θ denote the zero of either X or $[X]$ and let I be the identity in $[X]$. Let U and V be the solutions to the operator equation

$$Y'' = AY' + BY \quad (3)$$

satisfying the initial conditions $U(0) = \Theta$, $U'(0) = I$ and $V(0) = I$, $V'(0) = \Theta$. Since solutions to initial value problems for either (1) or (3) exist uniquely on any interval where A and B are continuous, the solution to (1) satisfying the initial conditions $y(0) = \alpha$, $y'(0) = \gamma$ is given by

$$y(t) = U(t)\gamma + V(t)\alpha. \quad (4)$$

Consequently, we have

LEMMA 1. *On a compact interval $[0, \tau]$, solutions to (1), (2) are unique if, and only if, $U(\tau)$ is one-one and (1), (2) is solvable for all α, β if, and only if, $U(\tau)$ is onto.*

LEMMA 2. *Let A and B be continuous on $[0, \tau]$ into $[X]_c$, the ideal in $[X]$ of completely continuous operators. Then $U(\tau)$ is one-one if, and only if, $U(\tau)$ is onto.*

Proof. Double integration of (3) together with the initial conditions yields $U(\tau) = \tau I + \int_0^\tau (\tau - s)[AU' + BU] ds$. If L denotes the integral term on the right, the conditions on A and B insure that $L \in [X]_c$. Using the terminology in [11], let λ and μ denote the dimensions of the null space and the defect subspace of $\tau I + L$. By Theorem 3.1 of [11], the index $\mu - \lambda$ of $\tau I + L$ equals the index of τI when L is completely continuous. But the index of the invertible operator τI is zero so $\mu = \lambda$ and $U(\tau)$ is one-one ($\lambda = 0$) if, and only if, $U(\tau)$ is onto ($\mu = 0$).

We note that if the boundary conditions (2) are replaced by $y'(0) = \alpha$, $y(\tau) = \beta$ we can replace U by V in Lemma's 1 and 2. A similar correspondence exists between the remaining possible boundary conditions and the operators U' and V' .

The hypothesis of complete continuity in Lemma 2 cannot be dropped. For let $X = l_2$ and for $n \geq 1$ let $e_n = (0, \dots, 0, 1, 0, \dots)$ denote the usual complete orthonormal set. Define the constant operator B by

$$Bx = \sum x_n B e_n = -\sum x_n [n\pi/(n+1)]^2 e_n \text{ for } x = \sum x_n e_n \in X.$$

B is linear, $\|B\| = \pi^2$ and the solution U to (3) is given in terms of the complete orthonormal set by

$$U(t)e_n = [(n+1)/n\pi] \sin [tn\pi/(n+1)] e_n.$$

Then $U(1)$ is one-one but $\Sigma (1/n)e_n$ does not belong to its range so $U(1)$ is not onto.

LEMMA 3. *If C is continuous on $[0, \tau]$ into $[X]$ and $Z(t)$ is a solution to $Z' = CZ$ in $[X]$ with $Z(0)$ invertible, then $Z(t)$ is invertible on $[0, \tau]$.*

Proof. On some interval to the right of zero, $Z^{-1}(t)$ exists and is a solution of $W' = -WC$. Applying Gronwall's inequality to $\|W(t)\| \leq \|W(0)\| + \int_0^t \|W(s)\| \|C(s)\| ds$ shows that $Z^{-1}(t)$ remains bounded on any subinterval of $[0, \tau]$ where it exists. Since the invertible operators form an open set in $[X]$, Z^{-1} must exist on $[0, \tau]$.

LEMMA 4. *For A continuously differentiable and B continuous on $[0, \tau]$ into $[X]$, equation (1) is equivalent to the system $z' = Cz$ under the substitution $y(t) = R(t)z(t)$ where $R(t)$ is the solution in $[X]$ of $R' = \frac{1}{2}AR$, $R(0) = I$ and $C(t) = R^{-1}(t)[\frac{1}{4}A^2(t) + B(t) - \frac{1}{2}A'(t)]R(t)$. If A and B are constant and commute, $C = \frac{1}{4}A^2 + B$.*

Proof. The first statement is a straight forward calculation. The last statement is a consequence of $R(t) = \exp(\frac{1}{2}tA)$ which commutes with A^2 and B when A and B commute.

3. CONSTANT COEFFICIENTS

Assume $A, B \in [X]$ are independent of $t \geq 0$ and $AB = BA$. Then the solution U to (3) is given by

$$U(t) = \sum_{n=0}^{\infty} \sum_{k=0}^n \frac{t^{n+k+1} n!}{(n+k+1)! (n-k)! k!} A^{n-k} B^k. \quad (5)$$

Moreover, $V(t) = U'(t) - AU(t)$ so that every solution of (1) can be written in terms of the above series using (4).

Define the complex valued function $u(t, \lambda, \mu)$ for t real, λ and μ complex numbers, by

$$u(t, \lambda, \mu) = \begin{cases} te^{\lambda(t/2)} & \lambda^2 - 4\mu = 0 \\ 2e^{\lambda(t/2)}(\lambda^2 + 4\mu)^{-1/2} \sinh(t/2)(\lambda^2 - 4\mu)^{1/2} & \lambda^2 - 4\mu \neq 0 \end{cases} \quad (5')$$

Then $u(t, \lambda, \mu)$ is the solution to the scalar equation

$$y'' = \lambda y' + \mu y \quad y(0) = 0, \quad y'(0) = 1$$

and has the expansion of (5) with λ, μ replacing A and B . The function u plays an important role in the following theorem.

THEOREM 1. *Let X be a complex Banach space and assume $A, B \in [X]$ commute. Then the BVP (1), (2) is uniquely solvable on $[0, \tau]$ if, and only if, $\tau(\lambda^2 + 4\mu)^{1/2} \neq 2n\pi i$ for all integers $n \neq 0$ and all pairs (λ, μ) belonging to the joint spectrum S of A, B .*

Proof. Let $C\{A, B\}$ denote the commutative subalgebra of $[X]$ generated by A and B . Let R be the ring of analytic complex valued functions $f(\lambda, \mu)$ defined on S and assume the non-equality holds. Then the function $u(\tau, \lambda, \mu)$ defined in (5') belongs to R and does not vanish on the compact set S . Since A and B are generators of $C\{A, B\}$, there is a homomorphism of R into $C\{A, B\}$ such that the functions $f(\lambda, \mu) = 1$, $g(\lambda, \mu) = \lambda$, $h(\lambda, \mu) = \mu$ in R correspond to I, A and B respectively in $C\{A, B\}$. ([12], pp. 91-94) From (5) and (5') we see that $U(\tau)$ corresponds to $u(\tau, \lambda, \mu)$ and since $u^{-1}(\tau, \lambda, \mu)$ belongs to R , the corresponding element in $C\{A, B\}$ must be $U^{-1}(\tau)$. It then follows from Lemma 1 that (1), (2) is uniquely solvable on $[0, \tau]$.

Suppose, conversely, that $\tau(\lambda^2 + 4\mu)^{1/2} = 2n\pi i$ for some $n \neq 0$, $(\lambda, \mu) \in S$. Then $u(\tau, \lambda, \mu)$ has a zero in S . But the spectrum of $U(\tau)$ is precisely the image of S under the mapping u , i.e. $\sigma(U(\tau)) = \{u(\tau, \lambda, \mu) : (\lambda, \mu) \in S\}$. ([12], p. 279) Consequently $0 \in \sigma(U(\tau))$ and $U(\tau)$ is not invertible so that either solutions to (1), (2) are not unique or do not exist for certain boundary values.

THEOREM 2. *If X is any Banach space and $A, B \in [X]$ are constant and commute with $\|\frac{1}{4}A^2 + B\| \tau^2 < \pi^2$, the BVP (1), (2) is uniquely solvable on $[0, \tau]$.*

Proof. According to Lemma's 1 and 4 it suffices to show that $U(\tau)$ is invertible where U is the solution in $[X]$ to $Z''(t) = CZ(t)$, $Z(0) = \Theta$, $Z'(0) = I$ for $C = \frac{1}{4}A^2 + B$.

If X is a complex space, the result follows from Theorem 1. For replacing A and B by Θ and C we need $\tau(4\mu)^{1/2} \neq 2n\pi i$ for $\mu \in \sigma(C)$, $n \neq 0$. But $\sigma(C) \subset \{\mu : |\mu| \leq \|C\|\} \subset \{\mu : |\mu| < (\pi/\tau)^2\}$ where the last inclusion follows from the hypothesis. Thus the non-equality holds for $n \neq 0$.

If X is not complex, $X^+ = X \times X$ with component wise addition and scalar multiplication defined by $(a + ib)(x, y) = (ax - by, bx + ay)$ is a complex Banach space with $\|(x, y)\|^+ = \max\{\|x\|, \|y\|\}$. For $C \in [X]$, $C^+(x, y) = (Cx, Cy) \in [X^+]$ is the extension of C . Since $\|C\| = \|C^+\|^+$, we can apply the argument of the above paragraph to X^+ and C^+ and assert that $U^-(\tau)$ is invertible. It follows that $U(\tau)$ is invertible on X .

The choice $A = \Theta$, $B = -I$ (so that $U(t) = \sin tI$) shows that π^2 cannot be replaced in the above inequality by any smaller constant.

The constant coefficient case with noncommuting A and B will be postponed until the next section. One expects that an inequality rather similar to the one given in Theorem 2 would be sufficient for (1), (2) to be uniquely solvable. However, the technique of reducing equation (1) to an equivalent system $z'' = Cz$ without the y' terms requires a non-constant operator $C(t) = \frac{1}{4}A^2 + R^{-1}(t)BR(t)$ and Theorem 1 is no longer applicable.

4. VARIABLE COEFFICIENTS

The next two theorems are obtained by considering a Fredholm integral equation equivalent to (1), (2). Conditions are put on A , B and $\tau > 0$ so that the integral operator is a contraction mapping on an appropriate function space.

THEOREM 3. *Let $A(t)$ be continuously differentiable and $B(t)$ continuous on $[0, \tau]$ into $[X]$ and let $C(t)$ be as in Lemma 4. If $\int_0^\tau s(\tau - s) \|C(s)\| ds < \tau$, the BVP (1), (2) is uniquely solvable.*

Proof. Let F denote the Banach space of continuous X -valued functions $u(t)$ defined on $[0, \tau]$ with $\|u\|_F = \sup\{\|u(t)\| : 0 \leq t \leq \tau\}$. For fixed $\alpha, \beta \in X$ define the mapping L on F into F by $L(u) = w$ where

$$w(t) = \int_0^\tau G(t, s) C(s) u(s) ds + \frac{t}{\tau} (\beta - \alpha) + \alpha$$

and where

$$G(t, s) = \begin{cases} s(t - \tau)/\tau & 0 \leq s \leq t \\ t(s - \tau)/\tau & t \leq s \leq \tau \end{cases}$$

is the Green's function for the scalar equation $y'' = 0$. Let $\lambda = \tau^{-1} \int_0^\tau s(\tau - s) \|C(s)\| ds < 1$. Then for $u, v \in F$

$$\|L(u - v)(t)\| \leq \int_0^\tau \|G(t, s)\| \|C(s)\| ds \|u - v\|_F \leq \lambda \|u - v\|_F$$

so $\|(L(u - v))\|_F \leq \lambda \|u - v\|_F$ and L is a contraction. The unique fixed point of L is the (unique) solution to $z'' = Cz$, $z(0) = \alpha$, $z(\tau) = \beta$. Since L is a contraction independent of the choice of α, β we have that the BVP for the reduced equation is uniquely solvable on $[0, \tau]$. By Lemma 4 the same is true for (1), (2).

COROLLARY 1. *If A is continuously differentiable and B is continuous on $[0, \tau]$ into a commutative subalgebra of $[X]$ with $\int_0^\tau s(\tau - s) \|\frac{1}{4}A^2(s) + B(s) - \frac{1}{2}A'(s)\| ds < \tau$, the BVP (1), (2) is uniquely solvable.*

Proof. Under the present hypotheses the operator $R(t)$ in Lemma 4 commutes with A^2 , B and A' on $[0, \tau]$ so that $C = \frac{1}{4}A^2 + B - \frac{1}{2}A'$ in Theorem 3.

As a special case of this corollary with $A = \Theta$ and X the real numbers we have; If a valued solution $u(t) \neq 0$ of $u'' = bu$ has two zeros in $[0, \tau]$ then $\int_0^\tau s(\tau - s) |b(s)| ds \geq \tau$. (cf Theorem 5.1, p. 345 of [2]).

COROLLARY 2. *If B is continuous on $[0, \tau]$ into $[X]$ and satisfies any of*

- (i) $\tau \int_0^\tau \|B(t)\| dt \leq 4$
- (ii) $\tau^2 \sup\{\|B(t)\| : 0 \leq t \leq \tau\} < 8$
- (iii) $\tau^3 \int_0^\tau \|B(t)\|^2 dt < 48$

then (1), (2) with $A = \Theta$ is uniquely solvable.

Proof. When $A = \Theta$ we have $C = B$ in Theorem 3. From the inequalities $s(\tau - s) \leq \tau^2/4$, $\int_0^\tau |G(t, s)| ds \leq \tau^2/8$ and $\int_0^\tau |G(t, s)|^2 ds \leq \tau^3/48$ we $\int_0^\tau |G(t, s)| \|B(s)\| ds < 1$ using (i), (ii), and (iii) respectively, where the Schwartz inequality is used for (iii). In each case the mapping of Theorem 3 is a contraction.

A restatement of condition (i) gives Lyapunov's result on the zeros of solutions to (1).

COROLLARY 3. [Lyapunov] *A necessary condition for the system $y'' = By$ to have a nontrivial solution with two zeros on $[0, \tau]$ is that $\int_0^\tau \|B(t)\| dt > 4/\tau$.*

When A and B are constant but $AB \neq BA$, we can replace $B(t)$ in Corollary 2 (or Theorem 3) by $C(t) = \frac{1}{4}A^2 + \exp((-t/2)A) B \exp((t/2)A)$

to obtain sufficient conditions for unique solvability. Weaker but more easily verified sufficient conditions are obtained by using the estimate $\|C(t)\| \leq \| \frac{1}{4}A^2 + B \| e^{t\|A\|}$ which follows by taking norms through the equation $C(t) = [\frac{1}{4}A^2 + B] + \frac{1}{2} \int_0^t [C(s)A - AC(s)] ds$ and applying Gronwall's inequality. Using this estimate, which is clearly valid when $AB = BA$, in Corollary 2 we get

COROLLARY 4. *If $A \neq \Theta$ and B are constant, then each of the following is sufficient that (1), (2) be uniquely solvable.*

- (i) $\tau \| \frac{1}{4}A^2 + B \| (e^{\tau\|A\|} - 1) \leq 4 \| A \|^2$
- (ii) $\tau^2 \| \frac{1}{4}A^2 + B \| e^{\tau\|A\|} < 8$
- (iii) $\tau^3 \| \frac{1}{4}A^2 + B \| (e^{2\tau\|A\|} - 1) < 96 \| A \|^3$

THEOREM 4. *Let $h(s) = \| A(s) \| + s \| B(s) \|$ and assume that $\max\{\int_0^\tau sh(s) ds, \int_0^\tau (\tau - s) h(s) ds\} < \tau$. Then (1), (2) is uniquely solvable on $[0, \tau]$.*

Proof. Let F denote the class of continuously differentiable X valued functions u defined on $[0, \tau]$ satisfying $u(0) = \Theta$. Then F with norm defined by $\|u\|_F = \sup\{\|u'(t)\| : 0 \leq t \leq \tau\}$ is a Banach space. Define the mapping L on F into F by $L(u) = w$ where

$$w(t) = \int_0^\tau G(t, s)[Au' + Bu] ds + \frac{t}{\tau} \beta$$

for $\beta \in X$ fixed. If the inequality of the theorem holds,

$$\lambda = \sup \left\{ \int_0^\tau |G_t(t, s)| h(s) ds : 0 \leq t \leq \tau \right\} < 1.$$

Since

$$\begin{aligned} \|L(u)'(t) - L(v)'(t)\| &= \left\| \int_0^\tau G_t(t, s) [A(u' - v')] \right. \\ &\quad \left. + B \int_0^s (u' - v') dr \right\| ds \leq \int_0^\tau |G_t(t, s)| h(s) ds \|u - v\|_F \\ &\leq \lambda \|u - v\|_F \end{aligned}$$

we have that L is a contraction. Again the argument is independent of the choice of β so (1), (2) with $\alpha = \Theta$ is uniquely solvable. But this means $U(\tau)$ in Lemma 1 is invertible, hence the conclusion.

COROLLARY. *If A and B are constant and $3\tau \|A\| + 2\tau^2 \|B\| < 6$, (1), (2) is uniquely solvable.*

THEOREM 5. Assume A, A' and B are continuous with $2 \int_0^t s \|A(s) + (t-s)[B(s) - A'(s)]\| ds \leq t$ on $[0, \tau]$. Then (1), (2) is uniquely solvable.

Proof. For the operator U of Lemma 1 we have $\|U(t) - tI\| < t$ on some interval $0 < t < \epsilon$. Let $(0, \lambda)$ denote the maximal interval on which the strict inequality holds. Then $\|U(t)\| < 2t$ on $(0, \lambda)$. If $\lambda \leq \tau$, we have

$$\begin{aligned} \lambda = \|U(\lambda) - \lambda I\| &= \left\| \int_0^\lambda [A + (\lambda - s)(B - A')] U ds \right\| \\ &< 2 \int_0^\lambda s \|A + (\lambda - s)(B - A')\| ds \leq \lambda, \end{aligned}$$

a contradiction. Hence $\lambda > \tau$ so that $\|\tau^{-1}U(\tau) - I\| < 1$ and $U(\tau)$ is invertible.

Unfortunately the preceding conditions are monotone in τ . When $X = [X] = \text{real numbers}$, the solution U of (3) is simply the scalar solution u to (1) satisfying $u(0) = 0$, $u'(0) = 1$. Since the zeros of u can only cluster at $+\infty$, (1), (2) is solvable on $[0, \tau]$ for almost every $\tau > 0$. In the general case one can construct an example of a system (1) such that the solution U of (3) has singularities (real numbers $t \geq 0$ for which $U(t)x = \theta$ for some $\theta \neq x \in X$) with a finite cluster point. However, it seems likely that in many cases $U(t)$ might again become invertible for t sufficiently large. If A is continuously differentiable and $K(t, s) = A(s) + (t-s)[B(s) - A'(s)]$ for $0 \leq s \leq t$, U solves the equation $U(t) = tI + \int_0^t K(t, s) U(s) ds$. Using successive iterations we can represent the solution formally as

$$U(t) = tI + \int_0^t sH(t, s) ds \quad \text{where} \quad H(t, s) = \sum_{n=0}^{\infty} K_n(t, s)$$

and

$$K_n(t, s) = \begin{cases} K(t, s) & n = 0 \\ \int_0^t K(t, u) K_{n-1}(u, s) du, & n \geq 1 \end{cases}$$

For $\tau > 0$ fixed, the series $H(t, s)$ converges in norm uniformly on $0 \leq s \leq t \leq \tau$. If

$$\left\| \int_0^\tau sH(\tau, s) ds \right\| < \tau \quad \text{for some} \quad \tau > 0, \quad \text{then} \quad (6)$$

$\|\tau^{-1}U(\tau) - I\| < 1$ and $U(\tau)$ is invertible. For certain specified A and B it may be possible to obtain estimates which imply (6) for certain large values of τ and hence (1), (2) would be solvable on such intervals.

5. HILBERT SPACE

In this section X will be a real Hilbert space with the inner product of two elements $x, y \in X$ denoted by $x \cdot y$. For an operator $T \in [X]$ we write T^* for the adjoint of T , T^0 for $\frac{1}{2}(T + T^*)$ and $T \geq 0$ in case $x \cdot Tx \geq 0$ for all $x \in X$.

The following condition is given in [3] as a sufficient condition that (1) be disconjugate on $[0, \tau]$. A minor modification of the original proof shows that the condition also implies that (1), (2) is solvable.

THEOREM 6. *In order that (1), (2) be uniquely solvable on $[0, \tau]$, it is sufficient that there exist a continuously differentiable operator $K(t)$ such that $B + K' \geq (\frac{1}{2}A + K^0)(\frac{1}{2}A^* + K^0)$ on $[0, \tau]$.*

Proof. Again we will show that $U(\tau)$ is invertible.

Let y be any solution of (1) such that $y(0) = \Theta$; i.e. $y(t) = U(t)x$ for some $x \in X$. Then for $0 < t \leq \tau$,

$$\begin{aligned} & y(t) \cdot y'(t) + y(t) \cdot K(t)y(t) \\ &= \int_0^t [(y \cdot y') + (y \cdot Ky)'] ds \\ &= \int_0^t [y' \cdot y' + y \cdot Ay' + y \cdot By + 2y' \cdot K^0y + y \cdot K'y] ds \\ &= \int_0^t [\|y' + (\frac{1}{2}A^* + K^0)y\|^2 + y \cdot Ly] ds \end{aligned} \quad (7)$$

where $L = B + K' - (\frac{1}{2}A + K^0)(\frac{1}{2}A^* + K^0) \geq 0$. If $y(\lambda) = \Theta$ for any $0 < \lambda \leq \tau$, the integrand on the right must be identically zero on $[0, \lambda]$ so y is a solution to the first order *IVP* $y' = -(\frac{1}{2}A^* + K^0)y$, $y(0) = \Theta$. By uniqueness we must have $y \equiv \Theta$ and it follows that $U(t)$ is one-one on $(0, \tau]$.

Now U is invertible on some open interval to the right of zero so if U fails to remain invertible on $[0, \tau]$ there is a first point λ , $0 < \lambda \leq \tau$ such that $\lim_{t \rightarrow \lambda^-} \|U^{-1}(t)\| = +\infty$. Since U is already one-one at λ , there must exist a sequence $\{x_n\}$ in X with $\|x_n\| = 1$ and $\lim_{n \rightarrow \infty} U(\lambda)x_n = \Theta$. But this is incompatible with (7). For the initial conditions $U(0) = \Theta$, $U'(0) = I$ require that $\|Z(t) - I\| \leq \frac{1}{2}$ on some interval $0 \leq t \leq \epsilon < \lambda$ where $Z = U' + (\frac{1}{2}A^* + K^0)U$. Consequently Z^{-1} exists and is bounded on $[0, \epsilon]$, say $\|Z^{-1}(t)\| \leq M$. It follows that $\|Z(t)x\| \geq (1/M)$ on $[0, \epsilon]$ for $x \in X$ with $\|x\| = 1$. Putting $y_n(t) = U(t)x_n$ into (7) we get

$$0 \geq \lim_{n \rightarrow \infty} \int_0^\lambda \|Z(t)x_n\|^2 ds \geq \frac{\epsilon}{M^2}.$$

From this contradiction we must have $U(\tau)$ invertible.

For $A = \Phi$, the inequality in Theorem 6 becomes $K' + B \geq K^0$ so (1), (2) is uniquely solvable on any interval $[0, \tau]$ on which the "Riccati" equation $K' + B = K^0$ has a solution.

COROLLARY. *Let A, A' and B be continuous on $[0, \tau]$ into $[X]$. Then (1), (2) is uniquely solvable if any of the following conditions hold on $[0, \tau]$.*

- (i) $A = A^*$ and $B \geq \frac{1}{2}A'$
- (ii) $A = -A^*$ and $B + \frac{1}{4}A^2 \geq \frac{1}{2}A'$
- (iii) $B \geq \frac{1}{4}AA^*$
- (iv) $B \geq A' + \frac{1}{4}A^*A$

Proof. In (i) and (ii) take $K = -\frac{1}{2}A$. Taking $K = \Theta$ and $-A$ gives (iii) and (iv).

From condition (i) we observe that for given B the system (1) can always be overdamped by taking $A(t) = g(t)I$ where $g(t)$ is any scalar function such that $g'(t) \leq -2 \|B(t)\|$ on $[0, \tau]$.

THEOREM 7. *Let X be a separable Hilbert space with $\{e_n\}$ a complete orthonormal set in X . Let $b_{ij}(t) = B(t) e_j \cdot e_i$, $a_{ij}(t) = A(t) e_j \cdot e_i$. If $a_{ij}(t) = 0$ and $b_{ij}(t) \leq 0$ on $[0, \tau]$ for $i \neq j$ while $\sum_{j=1}^{\infty} b_{ij}(t) \geq 0$ on $[0, \tau]$ for each $i \geq 1$. Then (1) is disconjugate on $[0, \tau]$. If A and B are also completely continuous operators, (1), (2) is uniquely solvable on $[0, \tau]$.*

Proof. It is convenient to think of A and B as infinite square matrices and y as an infinite column vector.

Suppose there is a nontrivial solution $y(t)$ to (1) with $y(t_1) = y(t_2) = \Theta$ for some $0 \leq t_1 < t_2 \leq \tau$. Taking y or $-y$, there is some component $y_k = y \cdot e_k$ of y and some $t_0 \in (t_1, t_2)$ such that

$$y_k(t_0) = \sup\{y_j(t) : t_1 \leq t \leq t_2, j \geq 1\} > 0.$$

Then $y'_k(t_0) = 0$ so

$$\begin{aligned} y''_k(t_0) &= a_{kk}(t_0) y'_k(t_0) + \sum_{j=1}^{\infty} b_{kj}(t_0) y_j(t_0) \\ &\geq \sum_{y_j(t_0) > 0} b_{kj}(t_0) y_j(t_0) \geq \left(\sum_{j=1}^{\tau} b_{kj}(t_0) \right) y_k(t_0) \geq 0 \end{aligned}$$

which contradicts y_k having a positive interior maximum. Hence (1) is disconjugate on $[0, \tau]$. It follows from Lemma 2 that (1), (2) is uniquely solvable with the additional assumption of complete continuity on A and B .

REFERENCES

1. HILLE, E. AND PHILLIPS, R., Functional analysis and semi-groups. *AMS Colloq. Publ.* **31** (1957).
2. HARTMAN, P., "Ordinary Differential Equations." Wiley, New York, 1964.
3. HARTMAN, P. AND WINTNER, A., On disconjugate differential systems. *Canad. J. Math.* **8** (1956), 72-81.
4. BARRETT, J. H., A Prufer transformation for matrix differential equations. *Proc. Am. Math. Soc.* **8** (1957), 510-518.
5. REID, W. T., A Prufer transformation for differential systems. *Pacific J. Math.* **8** (1958), 575-584.
6. REID, W. T., Oscillation criteria for self adjoint differential systems. *Trans. Am. Math. Soc.* **101** (1961), 91-106.
7. BARRETT, J. H., Matrix systems of second order differential equations. *Portugal. Math.* **14** (1955), 79-89.
8. STERNBERG, R. L., Variational methods and non-oscillation theorems for systems of differential equations. *Duke Math. J.* **19** (1952), 311-322.
9. ETGEN, G. J., Oscillatory properties of certain nonlinear matrix differential systems of second order. *Trans. Am. Math. Soc.* **122** (1966), 289-310.
10. HOWARD, H. C., Oscillation criteria for matrix differential equations. *Canad. J. Math.* **19** (1967), 184-199.
11. GOHBERG, I. AND KREIN, M., The basic propositions on defect numbers, root numbers and indices of linear operators. *AMS Transl. Ser.* **2**(13) (1960), 185-264.
12. GELFAND, I., RAIKOV, D., AND SHILOV, G., "Commutative Normed Rings." Chelsea, New York, 1964.